

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 September 2001 (07.09.2001)

PCT

(10) International Publication Number
WO 01/64676 A2

(51) International Patent Classification⁷: **C07D 471/10**,
235/06, 211/16, 401/06, 215/48, 401/14, 471/04, 405/12,
277/52, 209/08, 401/12, 403/12, 513/04, 473/06, 487/04,
409/06, 295/18, 211/28, 211/26, 401/10, 215/22, 307/68,
A61K 31/497, A61P 29/00

(21) International Application Number: PCT/US01/06715

(22) International Filing Date: 28 February 2001 (28.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/185,571 28 February 2000 (28.02.2000) US

(71) Applicant: SCIOS, INC. [US/US]; 820 West Maude Avenue, Sunnyvale, CA 94086 (US).

(72) Inventors: GOEHRING, Richard, R.; 6195 Potters Lane, Pipersville, PA 18947 (US). MAVUNKEL, Babu, J.; 1348 Avoset Terrace, Sunnyvale, CA 94087 (US). LIU, David, Y.; 201 Ferne Avenue, Palo Alto, CA 94306 (US). SCHREINER, George, F.; 12774 Leander Drive, Los Altos Hills, CA 94022 (US). LUEDTKE, Gregory; 963 La Mesa Terrace, #9, Sunnyvale, CA 94086 (US).

LEWICKI, John, A.; 308 Escobar Avenue, Los Gatos, CA 95030 (US).

(74) Agents: MURASHIGE, Kate, H. et al.; Morrison & Foerster LLP, 3811 Valley Centre Drive, Suite 500, San Diego, CA 92130-2332 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GI, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GI, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: INHIBITORS OF p38- α KINASE



(57) Abstract: The invention is directed to prepare medicaments for treatment of and to methods to treat conditions mediated by kinase using compounds of the formula (3) wherein Ar¹ and Ph are limited to specific embodiments or wherein the compound of formula (3) is a compound set forth in Figures 1A-II.

WO 01/64676 A2

INHIBITORS OF p38- α KINASE

Technical Field

The invention is directed to compounds that are useful in treating inflammation and that contain N-containing heterocycles such as piperazine or piperidine moieties coupled to phenyl and other aryl groups. More particularly, the invention concerns novel compounds of this type as well as methods to treat heart and kidney conditions using these and other compounds.

Background Art

A large number of chronic and acute conditions have been recognized to be associated with perturbation of the inflammatory response. A large number of cytokines participate in this response, including IL-1, IL-6, IL-8 and TNF. It appears that the activity of these cytokines in the regulation of inflammation rely at least in part on the activation of an enzyme on the cell signaling pathway, a member of the MAP kinase family generally known as p38 and alternatively known as CSBP and RK. This kinase is activated by dual phosphorylation after stimulation by physiochemical stress, treatment with lipopolysaccharides or with proinflammatory cytokines such as IL-1 and TNF. Therefore, inhibitors of the kinase activity of p38 are useful anti-inflammatory agents.

PCT applications WO98/06715, WO98/07425, WO98/28292 and WO 96/40143, all of which are incorporated herein by reference, describe the relationship of p38 kinase inhibitors with various disease states. As mentioned in these applications, inhibitors of p38 kinase are useful in treating a variety of diseases associated with chronic inflammation. These applications list rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions, sepsis, septic shock, endotoxic shock, Gram-negative sepsis, toxic shock syndrome, asthma, adult respiratory distress syndrome, stroke, reperfusion injury, CNS injuries such as neural trauma and ischemia, psoriasis, restenosis, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcosis, bone resorption diseases such as osteoporosis, graft-versus-host reaction, Crohn's Disease, ulcerative colitis including inflammatory bowel disease (IBD) and pyresis.

The above-referenced PCT applications disclose compounds which are p38 kinase inhibitors said to be useful in treating these disease states. These compounds are either imidazoles or are indoles substituted at the 3- or 4-position with a piperazine or piperidine ring linked through a carboxamide linkage. Additional compounds which are conjugates of piperazines with indoles are described as insecticides in WO97/26252, also incorporated herein by reference.


U.S. Patent No. 5,719,135 describes tyrosine kinase inhibitors containing piperidine or piperazine rings linked through a methylene at position 1 of piperidine to various aromatic systems which must further contain a γ lactam fused to a pyridine ring. Similar compounds are described in U.S. Patent No. 5,663,346 and in WO96/22976. Other cyclic tyrosine kinase inhibitors are described in PCT application WO95/06032. In addition, WO94/20062 describes balanoids as protein kinase C inhibitors. The balanoid compounds contain multiple aromatic systems which include at least a ring containing at least seven members. Some of the compounds useful in the method of the present invention are known compounds.

Additional compounds which contain piperidine or piperazine linked to an indole, benzimidazole, or benztriazole ring system are disclosed in PCT publication WO99/61426 published 2 December 1999; PCT publication WO00/59904 published 12 October 2000, and PCT publication WO00/_____ published _____ December 2000, the contents of which are incorporated herein by reference. Related compounds to those described herein are also set forth in PCT publication WO00/12074 published 9 March 2000 and incorporated herein by reference.

Disclosure of the Invention

The invention is directed to methods of treating inflammation generally, including specific conditions such as those described in the Background section above. The compounds of the invention have been found to inhibit p38 kinase and are thus useful in treating diseases mediated by this enzyme. These compounds also inhibit p38 α preferentially as compared to their inhibition of p38 β as is further discussed below.

The compounds useful in the invention are of the formula



and the pharmaceutically acceptable salts thereof, or a pharmaceutical composition thereof,

wherein


5 Z is N or CR¹, R¹ is a noninterfering substituent,

each of X¹ and X² is a linker,

Ar¹ and Ar² are identical or different, and represent optionally substituted C1-C20 hydrocarbyl residues wherein at least one of Ar¹ and Ar² is an optionally substituted aryl group, with the proviso that when X² is CH₂ or an isostere thereof, X¹ is CO or an isostere thereof, and Ar² is optionally substituted phenyl, Ar¹ is other than an optionally substituted indolyl, benzimidazolyl or benzotriazolyl substituent, and wherein said optionally substituted phenyl is not an optionally substituted indolyl, benzimidazolyl, or benzotriazolyl,

15 Y is a noninterfering substituent, wherein n is an integer from 0-4, and wherein m is an integer from 0-4 and l is an integer from 0-3.

Preferably, the compounds useful in the invention are of the formula



and the pharmaceutically acceptable salts thereof,

20 wherein Ar¹ is optionally substituted furanyl, thiophenyl, phenyl system having 0, 1, or 2 heterocyclic N atoms or naphthyl system having 0, 1, 2 or 3 heterocyclic N atoms;

X¹ is CO or an isostere thereof;

Y is optionally substituted alkyl (1-6C), optionally substituted aryl (6-10C), or optionally substituted arylalkyl (7-11C);

25 n is 0 or 1;


Z is CH or N;

X^2 is CH_2 or an isostere thereof; and

Ph is optionally substituted phenyl.

The optional substituents on the aryl moieties (including phenyl) include halo, nitro, optionally substituted alkyl (1-6C) or alkenyl (1-6C), CN, guanidino or CF_3 , as well as RCO, COOR, $CONR_2$, SO_2NR_2 , -OOCR, -NROCR, -NROCOR, NR_2 , OR or SR, wherein R is H or alkyl (1-6C), as well as substitution by phenyl, itself optionally substituted by the foregoing substituents. Any two substituents may form a 5-7 membered carbocyclic or heterocyclic ring subject to the proviso.

Thus, in one aspect, the invention is directed to compounds of the formulas set forth above and to pharmaceutical compositions containing them. In other aspects, the invention is directed to methods of treatment and preparation of medicaments using compounds of the formula set forth above. The invention is also directed to uses of specific classes of compounds within the genus of formula (1). In particular, the invention is related to compounds of formula (3)



In other aspects, the invention is directed to methods to produce the compounds useful in the invention.

In particular, the invention is directed to the use of compounds of formula (3) to prepare a medicament for treatment of conditions mediated by p38 α kinase or to methods of treatment of such conditions wherein the substituents in formula (3) are defined in the alternative as follows:

1. In formula (3), Ar^1 is a furanyl ring system optionally substituted by noninterfering substituents other than phenyl;

X^1 is CO or an isostere thereof;

25 Z is N or CH;

X^2 is CH_2 or an isostere thereof; and

Ph is optionally substituted phenyl.

2. In formula (3), Ar^1 is a thiophene ring system optionally substituted by a noninterfering substituent, and not fused to an additional ring;

30 X^1 is CO or an isostere thereof;

Z is N or CH;

X^2 is CH_2 or an isostere thereof; and

Ph is optionally substituted phenyl.

3. In formula (3), Ar^1 is an optionally substituted aryl group other than optionally substituted indolyl, benzimidazolyl, or benzotriazolyl;

5 X^1 is CO or an isostere thereof;

Z is N or CH;

X^2 is CH_2 or an isostere thereof; and

Ph represents optionally substituted phenyl

wherein at least one of Ar^1 and Ph is substituted by at least one substituent
10 comprising a saturated ring system optionally containing 1-2 N atoms and/or 1-2 O atoms.

4. The compound of formula (3) is a compound illustrated in Figures 1A-1I.

Brief Description of the Drawings

15 Figures 1A-1I show structures of additional compounds of the invention.

Modes of Carrying Out the Invention

The compounds of formula (1) set forth above are defined by the nature of the substituents on the heterocycloalkyl ring in the center of the formula. Piperazine or piperidine rings are preferred and piperidine rings are more preferred. The central
20 piperazine or piperidine ring can be expanded or contracted using $-CH_2$ groups so that it includes from 4 members up to 11 members. The substitution on this ring is on the N or Z positions only. Although not bound by this theory, the function of the central heterocycloalkyl group is apparently to space the Ar^2 group, which is generally hydrophobic, from the Ar^1 group, which is preferably but not necessarily hydrophilic.

25 The central heterocycloalkyl ring can include from 1-2 N. If the ring contains only 1 N then Z is a $-CR^1$ wherein R^1 is a noninterfering substituent. Preferably, R^1 is alkyl, alkoxy, aryl, arylalkyl, aryloxy, heteroaryl, halogen, acyl, carboxy, or hydroxy. More preferably, R^1 is hydroxy, alkyl, or alkoxy.

30 The linker that couples the central heterocycloalkyl ring and the Ar^1 and Ar^2 groups on either end of the molecule are preferably saturated or unsaturated alkylene

optionally containing 1-4 carbonyl, 1-4 SO₂, and/or 1-3 heteroatoms, including a linker which is CO, SO₂, SO or contains a heteroatom, and optionally substituted with a substituent selected from the group consisting of halo, alkyl, alkoxy, cycloalkyl, aryl, aryloxy, arylalkyl, haloalkyl, polyhaloalkyl, haloalkoxy, polyhaloalkoxy, cycloheteroalkyl, cycloheteroalkylalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, arylthio, arylsulfinyl, arylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, acyl, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, cyano, carboxy, hydroxy, tetrazolyl, imidazole, oxazole, triazole, and -SOR wherein R is hydroxy, alkyl, aryl, alkoxy, aryloxy, cycloalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, heteroarylalkoxy or cycloheteroalkylalkoxy.

Preferably the linker is a methylene group, a carbonyl group, a methylene group coupled to a carbonyl group, a methylene group having a methyl substituent or a methylene group containing an optionally substituted phenyl group. The alkylene group is C1-C8, and preferably C1-C4, and more preferably C1. Preferably, haloalkyl or polyhaloalkyl groups are CF₃ or CF₃CH₂ and haloalkoxy or polyhaloalkoxy groups are CF₃O or CF₃CH₂O.

Ar¹ or Ar² is an aryl group which is the residue of an aromatic hydrocarbon containing one or more rings optionally including one or more heteroatoms, selected from the group of O, N and S. Preferably the aryl group has 6-12 carbon atoms and up to 3 heteroatoms, and more preferably the aryl group has 6-8 carbon atoms and 1 or 2 heteroatoms. More preferably, the aryl group is a saturated or unsaturated 5-7 membered heterocycle and even more preferably a saturated or unsaturated 5-6 membered heterocycle. Most preferably, the aryl group is phenyl or residues of an optionally benzo-fused heterocycle containing up to 3 heteroatoms selected from S, N and O. The most preferable aryl groups are indolyl, isoquinolyl, quinolyl, benzimidazolyl, benzotriazolyl, benzothiazolyl, benzofuranyl, pyridyl, thienyl, furyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl, morpholinyl or piperidyl. When the linker is not CO or an isostere thereof, Ar¹ or Ar² is most preferably indolyl, benzimidazolyl or benzotriazolyl.

Ar¹ and Ar² and/or other aryl substituents are optionally substituted with substituents including one or more of halo, alkyl, alkoxy, cycloalkyl, aryl, aryloxy,

arylalkyl, haloalkyl, polyhaloalkyl, haloalkoxy, polyhaloalkoxy, cycloheteroalkyl, cycloheteroalkylalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, arylthio, arylsulfinyl, arylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, alkoxycarbonyl, aryloxy, heteroaryloxy, acyl, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, cyano, carboxy, hydroxy, tetrazolyl, imidazole, oxazole, triazole, and -SOR wherein R is hydroxy, alkyl, aryl, alkoxy, aryloxy, cycloalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, heteroarylalkoxy or cycloheteroalkylalkoxy.

Preferably the substituents include halo, alkoxy, alkoxyaryl, aminoalkyl, aminoaryl, and substituted aryl. Most preferably Ar^1 and Ar^2 include at least one halo, alkoxy, or N-containing substituent.

Y is selected from the group consisting of H, optionally substituted alkyl, alkoxy, cycloalkyl, cycloheteroalkyl, aryl, cycloheteroalkyl, heteroaryl, halogen, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, acyl, carboxy, hydroxy, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, cyano, amino, and alkylamino. Preferred groups are hydroxy, alkyl and carbonyl.

The compounds of formula (2) set forth above are defined by the nature of the substituents on the piperazine or piperidine ring. Piperidine rings are preferred.

In formula (2), Ar^1 is optionally substituted furanyl, thiophenyl, optionally substituted phenyl systems having 0, 1, or 2 heterocyclic N atoms or naphthyl systems having 0, 1, 2 or 3 heterocyclic N atoms. The nature of the substituents and the preferred substituents are discussed below.

X^1 is CO or an isostere thereof. Thus, in addition to CO, X^1 may be CH_2 , SO, SO_2 , or $CHOH$. CO is preferred.

Z is CH or N.

X^2 is CH_2 or an isostere thereof.

The aryl moieties represented by Ar^1 and Ph in formula (2) may optionally be substituted by substituents including straight or branched chain alkyl (1-6C), straight or branched chain alkenyl (2-6C), halo, RCO, COOR, $CONR_2$, SO_2NR_2 , -OOCR, -NROCR, -NROCOR, OR, SR, NR_2 , NO_2 , CN, or CF_3 , wherein R is H or straight or branched chain alkyl (1-6C). Phenyl moieties may also be substituted with an additional phenyl residue, preferably at the 4-position. Any two substituents may form a 5-7 membered carbocyclic or heterocyclic ring subject to the proviso. The

additional phenyl residue may itself be substituted with the substituents set forth above. The additional phenyl may be substituted in all five positions, but preferably less, preferably in 1-2 positions or not at all.

Preferred substituents include halo, alkyl (1-6C), OR, SR and NR₂, more preferably halo, OR and alkyl (1-4C), most preferably halo and OCH₃. The substituents on the phenyl moiety as an embodiment of Ar¹ or on Ph may occupy all five available positions, preferably 1-2 positions or the phenyl is unsubstituted. If Ar¹ comprises a pyridyl residue, only 4 positions are available; preferably only 1-2 positions are substituted or preferably the pyridyl is unsubstituted. If Ar¹ comprises furanyl or thiophenyl, only 3 positions are available; preferred numbers of substitutions in this case are 1 or 0.

n may be 0 or 1, and is preferably 0. However, when n is 1, Y is present and may be alkyl, arylalkyl or aryl, all of which may optionally be substituted by the substituents set forth above. Preferred embodiments of Y include unsubstituted alkyl (1-6C) and unsubstituted arylalkyl (7-11C), most preferably unsubstituted lower alkyl (1-4C).

In one group of compounds of formula (3), Ar¹ is a furanyl ring system optionally substituted by noninterfering substituents other than phenyl. These noninterfering substituents may include alkyl, alkoxy, cycloalkyl, cycloheteroalkyl, heteroaryl, halogen, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, acyl, carboxy, hydroxy, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, CF₃, cyano, amino and alkylamino.

Preferred substituents are alkyl, halo, CF₃, alkoxy, and cycloalkyl. Embodiments containing 0, 1 and 2 substituents are preferred, in particular embodiments with 0 or 1 substituents.

In a second embodiment of the compounds of formula (3), Ar¹ is a thiophene ring system containing optional substitutions with noninterfering substituents. The thiophene is not fused to an additional ring however. The optional substituents on the thiophene ring system may include alkyl, alkoxy, cycloalkyl, cycloheteroalkyl, heteroaryl, halogen, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, acyl, carboxy, hydroxy, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, CF₃, cyano, amino, alkylamino, and aryl. The preferred number of substituents is 0, 1 or 2, preferably 0 or 1.

In both the preceding embodiments - *i.e.*, wherein Ar¹ is a thiophene ring system unfused to another ring or a furanyl ring system, the phenyl group is optionally substituted by one or more noninterfering substituents. Preferably these substituents are alkyl, alkoxy, cycloalkyl, cycloheteroalkyl, heteroaryl, halogen, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, acyl, carboxy, hydroxy, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, CF₃, cyano, amino, alkylamino and aryl. Particularly preferred substituents are alkyl, alkoxy, and halo. It is preferred that the number of substituents be 0, 1 or 2, preferably 0 or 1 and more preferably 0.

In a third embodiment of the compounds of formula (3), although Ar¹ can be any optionally substituted aryl group other than optionally substituted indolyl, benzimidazolyl, or benzotriazolyl, it is required that at least one of Ar¹ and Ph is substituted by at least one substituent comprising a saturated ring system, optionally containing 1-2 N atoms or 1-2 O atoms. Preferred such substituents are cyclohexyl, cyclopentyl, piperazinyl, piperidinyl, or morpholinyl. It is preferred that only one saturated ring system be present as a substituent on Ar¹ or Ph. In addition to this required substituent, either or both of Ar¹ and Ph may contain additional substituents selected from those set forth in the previous embodiments as substituents on the phenyl moiety. Preferred substituents are alkyl, alkoxy and halo.

As set forth above, particularly preferred forms of the compounds of formula (3) are those set forth in figures 1A-1H.

The compounds of formula (1), (2) or (3) may be supplied in the form of their pharmaceutically acceptable acid-addition salts including salts of inorganic acids such as hydrochloric, sulfuric, hydrobromic, or phosphoric acid or salts of organic acids such as acetic, tartaric, succinic, benzoic, salicylic, and the like. If a carboxyl moiety is present on the compounds of formula (1), (2) or (3), the compound may also be supplied as a salt with a pharmaceutically acceptable cation.


In the event the compounds of formula (1), (2) or (3) contains one or more chiral centers, all of the stereoisomers are included within the scope of the invention, as well as mixtures thereof. Thus, the compounds of formula (1), (2) or (3) in these instances may be supplied as a single stereoisomer, as a racemic mixture, as a partially racemic mixture, or generally, a mixture of stereoisomers in any proportion.

The invention is directed, in addition to methods of treatment and of preparing medicaments, to compounds falling within the scope of formula (1), (2) or (3) as compositions of matter:

In one aspect, the invention is directed to compounds of the general formula (2), and the pharmaceutically acceptable salts thereof, wherein Ar¹ is optionally substituted furanyl, thiophenyl, or phenyl system containing 0, 1 or 2 N as heterocyclic atoms or naphthyl system containing 0, 1, 2 or 3 N as heterocyclic atoms, X¹ is CO or an isostere thereof; Y is optionally substituted alkyl (1-6C), optionally substituted aryl (6-10C), or optionally substituted arylalkyl (7-11C); n is 1; Z is N or CH; X² is CH₂ or an isostere thereof; and Ph is optionally substituted phenyl.

The invention is also directed to compounds of formula (2) and the pharmaceutically acceptable salts thereof, wherein Ar¹ is optionally substituted furanyl, thiophenyl, or phenyl system containing 1 or 2 N atoms as heterocyclic atoms or naphthyl system containing 0, 1, 2, or 3 N atoms as heterocyclic atoms, X¹ is CO or an isostere thereof; Y is optionally substituted alkyl (1-6C), optionally substituted aryl (6-10C), or optionally substituted arylalkyl (7-11C); n is 0-1; Z is N or CH; X² is CH₂ or an isostere thereof; and Ph is optionally substituted phenyl; wherein said substituents on Ar¹ and Ph are independently selected from the group consisting of straight or branched chain alkyl (1-6C), straight or branched chain alkenyl (2-6C), halo, RCO, COOR, CONR₂, SO₂NR₂, -OOCR, -NROCR, -NROCOR, OR, SR, NR₂, NO₂, CN, or CF₃, wherein R is H or straight or branched chain alkyl (1-6C).

The invention is also directed to a compound of the formula (2) and the pharmaceutically acceptable salts thereof, wherein Ar¹ is an ortho substituted furanyl, thiophenyl, or phenyl system containing 0, 1 or 2 N as heterocyclic atoms, or naphthyl system containing 0, 1, 2 or 3 N as heterocyclic atoms; said ortho substituent is straight or branched chain alkyl (1-6C), straight or branched chain alkenyl (2-6C), halo, RCO, COOR, CONR₂, SO₂NR₂, -OOCR, -NROCR, -NROCOR, OR, SR, NR₂, NO₂, CN, or CF₃, wherein R is H or straight or branched chain alkyl (1-6C), with the proviso that when said Ar¹ is phenyl and said ortho substituent is OR, either R must be alkyl (3-6C) or Ar¹ must be



where R ' is H, alkyl (1-6C), alkenyl (2-6C) or arylalkyl (7-12C);

X¹ is CO or an isostere thereof; Y is optionally substituted alkyl (1-6C),
optionally substituted aryl (6-10C), or optionally substituted arylalkyl (7-11C); n is 0-
5 1; Z is N or CH; and Ph is optionally substituted phenyl.

The ortho position is defined herein as the position in the ring adjacent the X¹
position, for example the 2-position in a 5-membered ring.

Preferred embodiments of the compounds of the invention are as described
above with respect to compounds useful in the invention methods. In particular
10 piperidine forms are preferred over piperazines.

Particularly preferred compounds of the invention include;





1-(2-methoxy-4-hydroxybenzoyl)-4-benzylpiperidine;






1-(2-methoxy-4-methoxybenzoyl)-4-benzylpiperidine;








1-(2-methoxy-4-benzyloxybenzoyl)-4-benzylpiperidine; and






15 1-(2-methoxy-4-methoxybenzoyl)-4-(4-fluorobenzyl)piperidine.






In addition, the following table illustrates preferred compounds of the
invention.








Compound #	Structure
1	
2	
3	
4	







5	
6	
7	
8	
9	






10	
11	
12	
13	
14	
15	
16	






17	
18	
19	
20	
21	








22	
23	
24	
25	
26	









27	
28	
29	
30	
31	
32	
33	









34	
35	
36	
37	
38	
39	









40	
41	
42	
43	
44	







45	 <chem>COc1ccc(cc1)C(=O)N2CCN(C2)C(=O)c3cc4c(cc3)nnc4</chem>
46	 <chem>FC(F)(F)c1ccc(cc1)/C=C/C(=O)N2CCN(C2)C(=O)c3cc4c(cc3)nnc4</chem>
47	 <chem>FC(F)(F)c1ccc(cc1)/C=C/C(=O)N2CCN(C2)C(=O)c3cc4c(cc3)nnc4</chem>
48	 <chem>c1ccccc1C2NCCN(C2)c3ccc(Cl)cc3C(=O)c4cc5c(cc4)nnc5</chem>
49	 <chem>Fc1ccc(cc1)C2NCCN(C2)c3ccc(F)cc3C(=O)c4cc5c(cc4)nnc5</chem>







50	
51	
52	
53	
54	
55	
56	







57	
58	
59	
60	
61	
62	
63	
64	







65	
66	
67	
68	
69	
70	
71	
72	







73	
74	
75	
76	
77	
78	
79	
80	


81	
82	
83	
84	
85	
86	

87	 <chem>COC1=CC=C(C(=C1)OC)C(=O)N2CCCCC2C(c3ccccc3)NC(=O)Cc4ccncc4</chem>
88	 <chem>COC1=CC=C(C(=C1)OC)C(=O)N2CCCCC2C(c3ccccc3)N</chem>
89	 <chem>COC1=CC=C(C(=C1)OC)C(=O)N2CCCCC2C(c3ccccc3)NC(=O)Cc4ccncc4</chem>
90	 <chem>c1ccc(cc1)n2c(=O)[nH]c3ccccc3n2C(=O)N4CCCCC4Cc5ccccc5</chem>
91	 <chem>COC1=CC=C(C(=C1)OC)C(=O)N2CCCCC2C(c3ccccc3)NC(=O)Cc4ccncc4</chem>
92	 <chem>COC1=CC=C(C(=C1)OC)C(=O)N2CCCCC2C(c3ccccc3)NC(=O)Cc4ccncc4</chem>

93	
94	
95	
96	
97	
98	

99	
100	
101	
102	
103	
104	

105	
106	
107	
108	
109	
110	



In addition to the compounds of formula (2) described above, the invention is directed to compounds of formula (3) as described above, and in particular the compounds of formula (3) set forth in figures 1A-1H.


All of the compounds of the invention are useful in methods to treat subjects wherein the subject harbors a condition which is mediated by an inappropriate p38 α kinase activity. Such conditions include various proinflammation conditions and various cardiac indications as described in the published documents cited above. The compounds of the invention are thus useful in preparing medicaments for the treatment of these conditions by admixing the compounds with suitable pharmaceutically acceptable excipients. Such excipients are well known in the art and are described in Remington's Pharmaceutical Sciences, latest edition, Mack Publishing Co., Easton, PA, incorporated herein by reference.

The invention is also directed to methods to prepare the compounds useful in the invention by forming the carboxamides from the appropriate aryl compounds as halides or free acids.


Synthesis Methods

Preferably, the compounds useful in the invention are synthesized, generally, by coupling an aryl moiety to a benzyl-substituted piperazine or piperidine. The general approach is shown in Reaction Schemes 1-6 hereinbelow. Reaction Scheme 3 is directed to the particular circumstance wherein Ar² includes an amino group to be functionalized in the presence of potentially competing groups.


Reaction Scheme 1




Reaction Scheme 2



Reaction Scheme 3




Reaction Scheme 4




5

Reaction Scheme 5



Reaction Scheme 6

5 Administration and Use

The methods of the invention are directed to treating conditions associated with p38 kinase activity, for example, inflammatory conditions. Thus, the compounds of formula (1) or their pharmaceutically acceptable salts are used in the manufacture of a medicament for prophylactic or therapeutic treatment of mammals, including humans, in respect of conditions characterized by excessive production of cytokines and/or inappropriate or unregulated cytokine activity on such cells as cardiomyocytes, cardiofibroblasts and macrophages.

The compounds of formulas (1) and (2) inhibit the production of cytokines such as TNF, IL-1, IL-6 and IL-8, cytokines that are important proinflammatory constituents in many different disease states and syndromes. Thus, inhibition of these cytokines has benefit in controlling and mitigating many diseases. The compounds of formulas (1) and (2) are shown herein to inhibit a member of the MAP kinase family variously called p38 MAPK (or p38), CSBP, or SAPK-2. The activation of this

protein has been shown to accompany exacerbation of the diseases in response to stress caused, for example, by treatment with lipopolysaccharides or cytokines such as TNF and IL-1. Inhibition of p38 activity, therefore, is predictive of the ability of a medicament to provide a beneficial effect in treating diseases such as coronary artery disease, congestive heart failure, cardiomyopathy, myocarditis, vasculitis, restenosis, such as occurs following coronary angioplasty, atherosclerosis, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions, multiple sclerosis, acute respiratory distress syndrome (ARDS), asthma, chronic obstructive pulmonary disease (COPD), silicosis, pulmonary sarcosis, sepsis, septic shock, endotoxic shock, toxic shock syndrome, heart and brain failure (stroke) that are characterized by ischemia and reperfusion injury, surgical procedures, such as transplantation procedures and graft rejections, cardiopulmonary bypass, coronary artery bypass graft, CNS injuries, including open and closed head trauma, inflammatory eye conditions such as conjunctivitis and uveitis, acute renal failure, glomerulonephritis, inflammatory bowel diseases, such as Crohn's disease or ulcerative colitis, graft vs host disease, bone resorption diseases like osteoporosis, type II diabetes, pyresis, psoriasis, cachexia, viral diseases such as those caused by HIV, CMV, and Herpes, and cerebral malaria.

Within the last several years, p38 has been shown to comprise a group of MAP kinases designated p38 α , p38 β , p38 γ and p38 δ . Jiang, Y. *et al. J Biol Chem* (1996) 271:17920-17926 first reported characterization of p38 β as a 372-amino acid protein closely related to p38 α . Kumar, S. *et al. Biochem Biophys Res Comm* (1997) 235:533-538 and Stein, B. *et al. J Biol Chem* (1997) 272:19509-19517 reported a second isoform of p38 β , p38 β 2 containing 364 amino acids with 73% identity to p38 α . All of these reports show evidence that p38 β is activated by proinflammatory cytokines and environmental stress, although the second reported p38 β isoform, p38 β 2, appears to be preferentially expressed in the CNS, heart and skeletal muscle compared to the more ubiquitous tissue expression of p38 α . Furthermore, activated transcription factor-2 (ATF-2) was observed to be a better substrate for p38 β 2 than for p38 α , thus suggesting that separate mechanisms of action may be associated with these forms. The physiological role of p38 β 1 has been called into question by the latter two reports since it cannot be found in human tissue and does not exhibit appreciable kinase activity with the substrates of p38 α .

The identification of p38 γ was reported by Li, Z. *et al. Biochem Biophys Res Comm* (1996) 228:334-340 and of p38 δ by Wang, X., *et al., J Biol Chem* (1997) 272:23668-23674 and by Kumar, S., *et al., Biochem Biophys Res Comm* (1997) 235:533-538. The data suggest that these two p38 isoforms (γ and δ) represent a
5 unique subset of the MAPK family based on their tissue expression patterns, substrate utilization, response to direct and indirect stimuli, and susceptibility to kinase inhibitors.

Various results with regard to response to drugs targeting the p38 family as between p38 α and either the putative p38 β 1 or p38 β 2 or both were reported by Jiang, Kumar, and Stein cited above as well as by Eysers, P.A. *et al. Chem and Biol* (1995) 5:321-328. An additional paper by Wang, Y. *et al. J Biol Chem* (1998) 273:2161-2168 suggests the significance of such differential effects. As pointed out by Wang, a number of stimuli, such as myocardial infarction, hypertension, valvular diseases, viral myocarditis, and dilated cardiomyopathy lead to an increase in cardiac workload
15 and elevated mechanical stress on cardiomyocytes. These are said to lead to an adaptive hypertrophic response which, if not controlled, has decidedly negative consequences. Wang cites previous studies which have shown that in ischemia reperfusion treated hearts, p38 MAPK activities are elevated in association with hypertrophy and programmed cell death. Wang shows in the cited paper that
20 activation of p38 β activity results in hypertrophy, whereas activation of p38 α activity leads to myocyte apoptosis. Thus, selective inhibition of p38 α activity as compared to p38 β activity will be of benefit in treating conditions associated with cardiac failure. These conditions include congestive heart failure, cardiomyopathy, myocarditis, vasculitis, vascular restenosis, valvular disease, conditions associated with
25 cardiopulmonary bypass, coronary artery bypass, grafts and vascular grafts. Further, to the extent that the α -isoform is toxic in other muscle cell types, α -selective inhibitors would be useful for conditions associated with cachexia attributed to TNF or other conditions such as cancer, infection, or autoimmune disease.

Thus, the invention is directed to the use of the compounds of formulas (1)
30 and (2) which selectively inhibit the activity of the p38 α isoform for treating conditions associated with activation of p38 α , in particular those associated with cardiac hypertrophy, ischemia or other environmental stress such as oxidation injury,

hyperosmolarity or other agents or factors that activate p38 α kinase, or cardiac failure, for example, congestive heart failure, cardiomyopathy and myocarditis.

5 The manner of administration and formulation of the compounds useful in the invention will depend on the nature of the condition, the severity of the condition, the particular subject to be treated, and the judgement of the practitioner; formulation will depend on mode of administration. As the compounds of formulas (1) and (2) are small molecules, they are conveniently administered by oral administration by compounding them with suitable pharmaceutical excipients so as to provide tablets, capsules, syrups, and the like. Suitable formulations for oral administration may also
10 include minor components such as buffers, flavoring agents and the like. Typically, the amount of active ingredient in the formulations will be in the range of 5%-95% of the total formulation, but wide variation is permitted depending on the carrier. Suitable carriers include sucrose, pectin, magnesium stearate, lactose, peanut oil, olive oil, water, and the like.

15 The compounds of formulas (1) and (2) may also be administered through suppositories or other transmucosal vehicles. Typically, such formulations will include excipients that facilitate the passage of the compound through the mucosa such as pharmaceutically acceptable detergents.

The compounds may also be administered topically, for topical conditions
20 such as psoriasis, or in formulation intended to penetrate the skin. These include lotions, creams, ointments and the like which can be formulated by known methods.

The compounds may also be administered by injection, including intravenous, intramuscular, subcutaneous or intraperitoneal injection. Typical formulations for such use are liquid formulations in isotonic vehicles such as Hank's solution or
25 Ringer's solution.

Alternative formulations include nasal sprays, liposomal formulations, slow-release formulations, and the like, as are known in the art.

Any suitable formulation may be used. A compendium of art-known formulations is found in Remington's Pharmaceutical Sciences, latest edition, Mack
30 Publishing Company, Easton, PA. Reference to this manual is routine in the art.

The dosages of the compounds of formula (1) will depend on a number of factors which will vary from patient to patient. However, it is believed that generally, the daily oral dosage will utilize 0.001-100 mg/kg total body weight, preferably from

0.01-50 mg/kg and more preferably about 0.01 mg/kg-10 mg/kg. The dose regimen will vary, however, depending on the conditions being treated and the judgment of the practitioner.


The inhibitors of p38 kinase can be used as single therapeutic agents or in combination with other therapeutic agents. Drugs that could be usefully combined with these compounds include natural or synthetic corticosteroids, particularly prednisone and its derivatives, monoclonal antibodies targeting cells of the immune system, antibodies or soluble receptors or receptor fusion proteins targeting immune or non-immune cytokines, and small molecule inhibitors of cell division, protein synthesis, or mRNA transcription or translation, or inhibitors of immune cell differentiation or activation.

As implicated above, although the compounds of formula (1) may be used in humans, they are also available for veterinary use in treating animal subjects.

The following examples are intended to illustrate but not to limit the invention.

Example 1

Preparation of 4-benzylpiperidin-1-yl benzene carboxamide (1-benzoyl-4-benzyl piperidine)



This example illustrates Reaction Scheme 1.

To a mixture of 4-benzylpiperidine (0.263 g, 1.5 mM) and diisopropylethylamine (0.53 mL, 3.0 mM) in 10 mL dichloromethane at room temperature was added benzoyl chloride (0.281 g, 2.0 mM) and the reaction mixture was stirred at room temperature for 20 h. The reaction mixture was poured into water and extracted with ethyl acetate (2 x 25 mL). The combined organic extract was washed with 1 N HCl, water and brine. The extract was dried over Na₂SO₄ and evaporated. The residue was chromatographed on a column of silica gel with ethyl acetate-hexane (10 to 50%,


gradient). Evaporation of the desired fractions gave 0.402 g (96%) of the title compound as an oil. ^1H NMR (CDCl_3) δ 7.10-7.40 (m, 10H), 4.60-4.70 (m, 1H), 3.70-3.80 (m, 1H), 2.80-2.90 (m, 1H), 2.70-2.80 (m, 1H), 2.60 (d, 2H), 1.70-1.80 (m, 1H), 1.50-1.70 (m, 2H), 1.20-1.40 (m, 2H). MS (ESI) m/e 279 (M^+)

5

Example 2

Preparation of 4-Benzylpiperidin-1-yl-(4-amino)benzene carboxamide (1-(4-aminobenzoyl)-4-benzyl piperidine)

This example illustrates Reaction Scheme 2:



10

4-Aminobenzoic acid (1.61g, 10 mM) was reacted with EDAC (1.92g, 10 mM) in 40 mL dry DMF for 15 minutes. 4-Benzylpiperidine (1.75g, 10 mM) was added followed by DMAP (20 mg, catalyst) and the reaction mixture was stirred for 20 h. The mixture was poured into water and extracted with methylene chloride (3x100 mL). The combined extract was washed with dilute hydrochloric acid, saturated sodium bicarbonate and water and dried over MgSO_4 . After evaporation of the solvent, the residue was chromatographed with methylene chloride-methanol (0 to 2% methanol, gradient) to yield 1.60 g (50%) of the title compound after recrystallization from ether-Hexane. ^1H NMR (CDCl_3): δ 7.40-7.10 (m, 7 H), 6.70 (d, 2 H), 3.95-3.90 (broad m, 2 H), 2.85-2.75 (broad m, 2 H), 2.6 (d, 2 H), 1.85-1.80 (m, 1 H), 2.75-2.65 (broad m, 2 H), 1.15-1.05 (broad m, 2 H). MS (ESI) m/e 294 (M^+).

15

20

Examples 3 and 4 illustrate Reaction Scheme 3:

Example 3

Preparation of 1-(2-chloro-4-aminobenzoyl)-4-benzyl piperidine

25

A. Preparation of the BOC-protected Intermediate

To a solution of 2-Chloro-4-*t*-butoxycarbonylaminobenzoic acid (0.50 g, 1.85 mM) in CH₂Cl₂ (8.0 mL) was added triethylamine (0.37 g, 0.52 mL, 3.70 mM) and 4-benzylpiperidine (0.65 g, 0.65 mL, 3.70 mM). This was followed by the dropwise addition of thionyl chloride (0.44 g, 0.27 mL, 3.70 mM). The mixture was allowed to stir at room temperature for 1 h, whereupon it was poured into water and extracted with CH₂Cl₂ (3 x 10 mL). The combined organics were dried over Na₂SO₄ and evaporated. Silica gel chromatography (5% methanol in dichloromethane) afforded 0.30 g (38%) of BOC-protected intermediate, i.e., 1-(2-chloro-4-*t*-butoxycarbonyl aminobenzoyl)-4-benzyl piperidine. ¹H NMR (CDCl₃): δ 7.30-7.10 (m, 2 H), 6.60 (s, 1 H), 4.70 (m, 1 H), 3.30 (m, 1 H), 2.50 (d, 2 H), 1.80 (m, 2 H), 1.60 (m, 2 H), 1.50 (s, 9 H), 1.20 (m, 3 H). MS (ESI) m/e 428 (M⁺).

B. 1-(2-chloro-4-*t*-butoxycarbonyl aminobenzoyl)-4-benzyl piperidine (0.15 g, 0.35 mM) from paragraph A was added to 4.0 N HCl in dioxane (2.0 mL). The mixture was stirred for 2 hours at room temperature whereupon it was triturated with Et₂O. The Et₂O was decanted and the residue was dried in vacuo to yield 0.10 g (87%) of the title compound. ¹H NMR (CDCl₃): δ 7.50 (m, 3 H), 7.30 (m, 4 H), 7.20 (d, 1 H), 7.10 (d, 2 H), 4.6 (t, 1 H), 3.30 (d, 1 H), 2.90 (t, 1 H), 2.70 (t, 1 H), 2.50 (s, 2 H), 1.70 (d, 2 H), 1.60 (t, 1 H), 1.20 (broad s, 2 H). MS (ESI) m/e 327 (M⁺-1).

Example 4

Preparation of 1-(2-chloro-4-methylaminobenzoyl)-4-benzyl piperidine

A solution of 1-(2-chloro-4-*t*-butoxycarbonyl aminobenzoyl)-4-benzyl piperidine (0.88g, 2.06 mM) in THF (4.0 mL) from paragraph A of Example 3 was added dropwise to an ice-cooled suspension of NaH in THF (4.0 mL). Stirring was continued for 0.5 hours before adding iodomethane (0.35 g, 0.15 mL, 2.47 mM) dropwise. The mixture was then allowed to stir at room temperature for 16 hours, at which time it was poured into water and extracted with ethyl acetate (3 x 10 mL). The combined extracts were dried over Na₂SO₄ and concentrated. The residue was dissolved in 4.0 N HCl in dioxane (10.0 mL) and stirred at room temperature for 2 hours whereupon it was triturated with Et₂O. The Et₂O was decanted and the residue was dried in vacuo to yield 0.56 g (79%) of 1-(2-chloro-4-methylaminobenzoyl)-4-benzyl piperidine. ¹H NMR (CDCl₃): δ 7.60-7.10 (m, 8 H), 4.70 (t, 1 H), 3.40 (

broad s, 1 H), 3.00 (m, 1 H), 2.90 (s, 3 H), 2.70 (m, 1 H) 2.60 (m, 1 H), 1.80 (broad s, 2 H), 1.60 (broad s, 1 H), 1.30 (m, 2 H). MS (ESI) m/e 342 (M^+).

Example 5

Preparation of Additional Compounds of Formula (2)

5 Using the techniques set forth in Examples 1-4, the following additional compounds of the invention were prepared:

1-(4-methoxybenzoyl)-4-benzyl piperidine,
1-(4-cyanobenzoyl)-4-benzyl piperidine,
1-(4-phenylbenzoyl)-4-benzyl piperidine,
10 1-(3-methoxybenzoyl)-4-benzyl piperidine,
1-(3,5-dimethoxybenzoyl)-4-benzyl piperidine,
1-(2-chlorobenzoyl)-4-benzyl piperidine,
1-(2-bromobenzoyl)-4-benzyl piperidine,
1-(2-iodobenzoyl)-4-benzyl piperidine,
15 1-(2-methoxybenzoyl)-4-benzyl piperidine,
1-(2-methylbenzoyl)-4-benzyl piperidine,
1-(2-phenylbenzoyl)-4-benzyl piperidine,
1-(2-(2-carbomethoxyethen-1-yl)benzoyl)-4-benzyl piperidine,
1-(2-naphthoyl)-4-benzyl piperidine,
20 1-(1-naphthoyl)-4-benzyl piperidine,
1-(2-furanoyl)-4-benzyl piperidine,
1-(2-thiophenoyl)-4-benzyl piperidine,
1-(2-dimethylaminobenzoyl)-4-benzyl piperidine,
1-(3-dimethylaminobenzoyl)-4-benzyl piperidine,
25 1-(4-dimethylaminobenzoyl)-4-benzyl piperidine,
1-(3,4,5-trimethoxybenzoyl)-4-benzyl piperidine,
1-(2-acetamidobenzoyl)-4-benzyl piperidine,
1-(3-acetamidobenzoyl)-4-benzyl piperidine,
1-(4-acetamidobenzoyl)-4-benzyl piperidine, and
30 1-(4-formamidobenzoyl)-4-benzyl piperidine,

Example 6

Preparation of 1-Phenyl-8-1,3,8-triazaspiro[4.5]decan-4-one-3-acetyl-4-benzylpiperidine

This example illustrates Reaction Scheme 4.

5 Acid I [*J Med Chem* (1996), 39(16):3169-3173]: 1.0 g, 2.57 mmol and
4-Benzylpiperidine, (0.46 g, 2.6 mmol) were treated with 0.51 g EDAC.HCl in 10 mL
dichloromethane (DCM) in the presence of 200 μ L 0.1M DMAP in DMF. After
stirring at room temperature for 4 hours, the reaction mixture was concentrated and
the residue was taken up in 50 mL ethyl acetate (EtOAc). The EtOAc layer was
10 washed with 10% aqueous sodium carbonate, 10% aqueous hydrochloric acid, sodium
chloride solution and dried over anhydrous sodium sulfate. Concentration gave crude
product as a white solid. This material was chromatographed using
ethylacetate/hexane on silica gel to give Boc-protected compound as a white solid.
700 mg EIMS M^+ 546. 100 mg of the Boc-protected compound was treated with 10
15 mL 4N HCl in dioxane for 60 mins. The reaction mixture was concentrated and the
solid obtained was filtered through a sintered glass funnel and washed with hexane.
Extensive drying gave the hydrochloride salt of Example 1. EIMS M^+ 446. 65 mg.

Example 7

Preparation of 4-Phenyl-3-pyrrolyl-4-benzylpiperidine

20 This example illustrates Reaction Scheme 4.

4-Phenyl-3-pyrrole carboxylic acid (333 mg, 3mmole) and 4-Benzylpiperidine
(525mg 3mmole) were dissolved in 15ml dichloromethane/dimethylformamide(1.1).
EDAC-HCl (573mg, 3mmole) was added and the reaction mixture was stirred
overnight. The solvent was removed and residue redissolved in EtOAc, the EtOAc
25 was washed with 10% HCl, 0.1M NaOH, water, the EtOAc extract was dried over
magnesium sulfate (anh). Evaporated EtOAc to obtain 489mg product.

Unless otherwise indicated all compounds were prepared following the
method outlined above using carboxylic acids that are either commercially available
or are available by synthesis, using methods known in the literature.

Example 8Preparation of (V) 4-(4')-Nitrophenyl-3-purroyl-4-benzylpiperidine

This example illustrates Reaction Scheme 6.

Preparation of II: Sodium hydride (614mg 25.6mmole) was suspended in
5 40ml diethyl ether. Methyl 4-nitrophenyl-trans-cinnamate (4.14g, 20mmole),
tosylmethylisocyanate (4.29g, 22mmole) dissolved in Et₂O/DMSO (2:1) 40ml were
added dropwise under nitrogen over 20min period. The mixture was stirred for 1 hour
at room temperature. Water was added to the reaction mixture and precipitate was
filtered. The aqueous layer was extracted with Et₂O (2 x 75ml), The combined
10 extracts were dried over MgSO₄ and solvent removed under reduced pressure to give
1.42g product. EIMS M⁺ 247.

Preparation of III: Methyl-3-(4-nitrophenyl)-2-pyrrole carboxylate (626mg
2.54mmole) was treated with potassium hydroxide (713mg, 12.7mmole) in
methanol/water (50:50). The reaction mixture was refluxed 2.5 hours. Methanol was
15 evaporated, the solution was cooled and acidified with HCl(aq) to pH 7. The mixture
was extracted with ethyl acetate, combined extracts dried over MgSO₄(anh.) and
concentrated to give 440mg product. EIMS M⁺ 232.

Preparation of IV: 3-(4-nitrophenyl)-2-pyrrole carboxylic acid (440mg
1.89mmole) and 4-Benzylpiperidine (331mg 1.89mmole) were dissolved in
20 DCM/DMF(10:1), EDAC (360mg, 1.89mmole) was added and the reaction mixture
was stirred for 18hr at room temperature. The solvent was evaporated, residue
dissolved in EtOAc, washed with 10% HCl, 0.10M NaOH, water, the EtOAc layer
was dried over MgSO₄, solvent evaporated to give 240mg product. Product was
chromatographed on silica gel with EtOAc:hex (8:2) to give 180mg purified product.
25 EIMS M⁺ 389.

Preparation of (V) 4-(4')-Nitrophenyl-3-purroyl-4-benzylpiperidine: 180mg of
the nitro compound was dissolved in 10ml methanol, 10% Pd/C (60mg) was added
and hydrogenated under hydrogen balloon for 2 hours. The catalyst was filtered over
celite, solvent was evaporated to give 136mg of the desired product. EIMS M⁺ 359.

Example 9

Assay for p38 Kinase Inhibition

The compounds to be tested were solubilized in DMSO and diluted into water to the desired concentrations. The p38 kinase was diluted to 10 $\mu\text{g/ml}$ into a buffer containing 20 mM MOPS, pH 7.0, 25 mM beta-glycerol phosphate, 2 mg/ml gelatin, 0.5 mM EGTA, and 4 mM DTT.

The reaction was carried out by mixing 20 μl test compound with 10 μl of a substrate cocktail containing 500 $\mu\text{g/ml}$ peptide substrate and 0.2 mM ATP (+200 $\mu\text{Ci/ml}$ gamma-32P-ATP) in a 4x assay buffer. The reaction was initiated by the addition of 10 μl of p38-kinase. Final assay conditions were 25 mM MOPS, pH 7.0, 26.25 mM beta-glycerol phosphate, 80 mM KCl, 22 mM MgCl_2 , 3 mM MgSO_4 , 1 mg/ml gelatin, 0.625 mM EGTA, 1 mM DTT, 125 $\mu\text{g/ml}$ peptide substrate, 50 μM ATP, and 2.5 $\mu\text{g/ml}$ enzyme. After a 40 minute incubation at room temperature, the reaction was stopped by the addition of 10 μl per reaction of 0.25 M phosphoric acid.

A portion of the reaction was spotted onto a disk of P81 phosphocellulose paper, the filters were dried for 2 minutes and then washed 4x in 75 mM H_3PO_4 . The filters were rinsed briefly in 95% ethanol, dried, then placed in scintillation vials with liquid scintillation cocktail.

Alternatively, the substrate is previously biotinylated and the resulting reactions are spotted on SAM²™ streptavidin filter squares (Promega). The filters are washed 4x in 2M NaCl, 4x in 2M NaCl with 1% phosphoric acid, 2x in water, and briefly in 95% ethanol. The filter squares are dried and placed in scintillation vials with liquid scintillation cocktail.

Counts incorporated are determined on a scintillation counter. Relative enzyme activity is calculated by subtracting background counts (counts measured in the absence of enzyme) from each result, and comparing the resulting counts to those obtained in the absence of inhibitor. IC_{50} values were determined with curve-fitting plots available with common software packages. Approximate IC_{50} values were calculated using formula

$$\text{IC}_{50} (\text{app}) = A \times i / (1 - A)$$

where A=fractional activity and i=total inhibitor concentration.


The percent inhibition of p38- α kinase was determined at various concentrations in order to determine IC_{50} values. Table 1 shows compounds tested for

the percent inhibition of p38- α kinase at concentrations of 15 μ M, 5 μ M, 1 μ M and 0.2 μ M. The table shows various embodiments of Ar¹X¹; unless otherwise noted, Ph is unsubstituted phenyl, Z is CH and n is 0. Thus, all of the compounds in Table 1 are 4-benzyl piperidines, unless otherwise noted. All show substantial inhibition at 15 μ M, some as high as 99%. Virtually all are inhibitory at 0.2 μ M.


Table 1	
Comp. No.	Ar ¹ X ¹
1	benzoyl
2	4-methoxybenzoyl
3	4-cyanobenzoyl
4	4-phenylbenzoyl
5	3-methoxybenzoyl
6	3,5-dimethoxybenzoyl
7	2-chlorobenzoyl
8	2-bromobenzoyl
9	2-iodobenzoyl
10	2-methoxybenzoyl
11	2-methylbenzoyl
12	2-(2-carbomethoxy ethen-1-yl) benzoyl
13	2-naphthoyl
14	1-naphthoyl
15	2-furanoyl
16	2-thiophenoyl
17	note 1
18	2-dimethylaminobenzoyl
19	3-dimethylaminobenzoyl
20	4-dimethylaminobenzoyl
21	2-pyridoyl
22	3-pyridoyl
23	4-pyridoyl
24	note 2
25	3,4,5-trimethoxybenzoyl
26	2-chloro-4-acetamidobenzoyl
27	2-4-dimethylbenzoyl
28	4-acetamidobenzoyl
29	4-formamidobenzoyl
30	2-aminobenzoyl
31	3-aminobenzoyl
32	4-aminobenzoyl
33	4,5-dimethoxybenzoyl

Table 1	
Comp. No.	Ar ¹ X ¹
34	2,4-dimethoxybenzoyl
35	2-hydroxybenzoyl
36	2-benzoyloxybenzoyl
37	3,5-bis-trifluoromethylbenzoyl
38	2,6-dimethoxybenzoyl
39	2-methylaminobenzoyl
40	6-quinolyl
41	2-chloro-4-benzamidoyl
42	2-hydroxy-4-aminobenzoyl
43	2-chloro-4-aminobenzoyl
44	2-chloro-4-methylaminobenzoyl
45	2-chloro-4-dimethylaminobenzoyl
46	2-methoxy-4-nitrobenzoyl
47	2-methoxy-4-aminobenzoyl
48	4-guanidinobenzoyl
49	2-bromobenzoyl (note 3)
50	phenylsulfonyl
51	4-amino-3-pyridoyl
52	2-methoxybenzoyl (note 4)

Note 1: In compound 17, X¹ is CO and Ar¹ has the structure:



Note 2: In compound 24, X¹ is CO and Ar¹ has the structure:



Note 3: In compound 49, n is 1 and Y is methyl. The methyl and benzyl group at the 4-position of the piperidine ring are *cis* to each other.

Note 4: In compound 52, Z is N; compound 52 is a piperazine derivative.

In some cases, IC₅₀ values have been determined as well as the ratio of IC₅₀ values for inhibition of β as compared to α p38 kinase. The compounds in Table 2 show IC₅₀ of about 200 nM to 1.5 μ M. Those tested are specific for p38- α at least by a factor of 5.

Table 2	
Comp. No.	Ar ¹ X ¹
1	benzoyl
2	4-methoxybenzoyl


Table 2	
Comp. No.	Ar ¹ X ¹
5	3-methoxybenzoyl
7	2-chlorobenzoyl
8	2-bromobenzoyl
9	2-iodobenzoyl
10	2-methoxybenzoyl
11	2-methylbenzoyl
16	2-thiophenoyl
18	2-dimethylaminobenzoyl
19	3-dimethylaminobenzoyl
20	4-dimethylaminobenzoyl
28	4-acetamidobenzoyl
29	4-formamidobenzoyl
32	4-aminobenzoyl
34	2,4-dimethoxybenzoyl
35	2-hydroxybenzoyl
38	2,6-dimethoxybenzoyl
39	2-methylaminobenzoyl
43	2-chloro-4-aminobenzoyl
44	2-chloro-4-methylaminobenzoyl
45	2-chloro-4-dimethylaminobenzoyl
49	2-bromobenzoyl (note 3)

Example 10

Preparation of Ortho-Substituted Aroyl Derivatives

The aroyl piperidines or piperazines of formula (2) can be provided with
 5 ortho-substituents via metallation as described by Beak, P. *et al.*, *J Org Chem* (1982) 47:34-46; Bindal, R.D. *et al.*, *J Org Chem* (1987) 52:3181-3185; Gshwend, H. *et al.*, *Org Rxns* (1979) 26:1. The general procedure is shown as follows:


Reaction Scheme 7



The starting material is first metallated with butyl lithium in the presence of base and then treated with an electrophile to provide the appropriate substitution. The electrophile may be an alkyl group, but additional functionality may be introduced by appropriate choice of the electrophile. For example, use of benzaldehyde as the

Compounds with further substitutions on the aroyl moiety can also be used as substrates, but those that contain an N-H require two equivalents of the butyl lithium/base.

Example 11A


Preparation of 4-benzylpiperidin-1-yl-(2-methoxy-4-benzyloxy)benzene carboxamide

A. Methyl 2-hydroxy-4-benzyloxybenzoate: To a solution of methyl 2,4-dihydroxybenzoate (17.5 g, 104 mM) in acetone at 0 °C were added potassium carbonate (52.5 g, 350 mM) and benzyl bromide (13.8 mL, 116 mM). The mixture was stirred at 0 °C for 2 h, followed by stirring at RT for 16 h. This mixture was filtered through a pad of Celite and concentrated. The resulting material was partitioned between ethyl acetate and water. The organic layer was washed with a

saturated sodium bicarbonate solution, dried (Na_2SO_4), filtered and concentrated. The residue was purified via silica gel chromatography with ethyl acetate-hexane (1 to 20 %, gradient). Evaporation of the desired fractions resulted in 13.48 g (50 %) of the title compound as a white solid. $^1\text{H-NMR}$ (CDCl_3) δ 7.80 (d, 1 H), 7.50-7.30 (m, 5 H), 6.55 (m, 2 H), 5.10 (s, 2 H), 3.95 (s, 3 H). MS (ESI) m/e 258 (m^+).

B. Methyl 2-methoxy-4-benzyloxybenzoate: To an ice-cooled solution of methyl 2-hydroxy-4-benzyloxybenzoate from paragraph A (12.92 g, 50.1 mM) in DMF (200 mL) was added NaH in a 60 % oil dispersion (3.00 g, 75.0 mM). The mixture was stirred for 5 min, at which time MeI (7.9 mL, 127 mM) was added dropwise. The mixture was allowed to stir at RT for 16 h, whereupon it was poured onto ice and extracted with ethyl acetate. The organics were washed with water and brine, then dried (Na_2SO_4) and concentrated to yield 13.11 g (96.3 %) of the title compound as a tan solid. $^1\text{H-NMR}$ (CDCl_3) δ 7.85 (d, 1 H), 7.50-7.30 (m, 5 H), 6.55 (m, 2 H), 5.15 (s, 2 H), 3.95 (s, 3 H), 3.90 (s, 3 H). MS (ESI) m/e 272 (m^+).

C. Title Compound: To a solution of 4-benzylpiperidine (27.5 mM, 4.85 mL) in toluene (75 mL) at -78°C was added Me_3Al (2 M solution in hexane, 32.5 mM, 16.25 mL). The mixture was stirred at -78°C for 10 min followed by stirring at RT for 1 h. The mixture was once again cooled to -78°C and a solution of methyl 2-methoxy-4-benzyloxybenzoate from paragraph B (20.0 mM, 6.96 g) in toluene (25 mL) was added dropwise. Stirring was continued at -78°C for 10 min, at which time the mixture was allowed to warm to RT and then refluxed for 16 h. The mixture was then cooled and quenched with water, made basic (pH 12) with 1 M NaOH, and stirred at RT for 0.5 h. The mixture was then extracted with ethyl acetate, washed with brine, dried (Na_2SO_4), filtered, and concentrated. The residue was purified via silica gel chromatography with ethyl acetate-hexane (10 to 60 %). Evaporation of the desired fractions resulted in 639 mg (74.8 %) of the title compound as an oil. $^1\text{H-NMR}$ (CDCl_3) δ 7.60-7.10 (m, 11 H), 6.60 (m, 2 H), 5.10 (s, 2 H), 4.75 (br d, 1 H), 3.80 (app d, 3 H), 3.55 (br d, 1 H), 3.00-2.50 (m, 4 H), 1.80 (m, 2 H), 1.60 (m, 1H), 1.30 (m, 1 H), 1.10 (m, 1 H). MS (ESI) m/e 414 (m^+).

Example 11BPreparation of 4-benzylpiperidin-1-yl-(2-methoxy-4-hydroxy)benzene carboxamide

A Parr hydrogenation vessel was charged with 4-benzylpiperidin-1-yl-(2-methoxy-4-benzyloxy)benzene carboxamide from Example 11A (15.00 g, 36.14 mM), dissolved in MeOH (100 mL), and 5 % palladium on carbon (1.2 g). The vessel was flushed, placed under 42 psi of H₂, and shaken for 16 h. The slurry was then filtered through Celite and concentrated. The crude material was recrystallized from ethyl acetate/hexane (1 : 10) to yield 11.24 g (95.7 %) of the title compound as a white solid. ¹H-NMR (CDCl₃) δ 7.40-6.80 (m, 6 H), 6.30 (m, 2 H), 4.75 (broad d, 2 H), 3.50 (app d, 3 H), 3.0-2.50 (m, 4 H), 1.75 (m, 2 H), 1.55 (m, 1 H), 1.20 (m, 1 H), 1.05 (m, 1 H). MS (ESI) *m/e* 324 (*m*⁺).

Example 12Effect of Ortho-Substitution

Using the methods set forth in Example 10, 1-benzoyl-4-benzylpiperidine isomers, substituted on the benzoyl substituent at various positions, were tested for their ability to inhibit p38α kinase at 15 μM concentration. The results are shown in Table 3:

Table 3: % Inhibition of p38α kinase at 15 μM			
	<i>ortho</i>	<i>meta</i>	<i>para</i>
F	98	92	94
Cl	99	80	85
Br	99	74	96
methyl	98	90	97
CF ₃	98	49	66

As shown in Table 3, for any given substituent, the ortho isomer shows enhanced inhibition.

The IC₅₀ values, obtained as set forth in Example 10, were also determined for the ortho-, meta-, and para-methoxy substituted benzoyl forms of 1-benzoyl-4-benzylpiperidine. These results are shown in Table 4:

Table 4: IC ₅₀ 's of methoxy analogs	
	IC ₅₀ (μM)
<i>ortho</i>	0.287
<i>meta</i>	1.1
<i>Para</i>	0.605

5

It is clear from these results that substitution at the ortho position is preferred.

The molar refraction values for the ortho derivatives containing chloro, bromo and methoxy substituents were determined as a measure of the size of the substituent as described by Hansch, C. *et al.*, *J Med Chem* (1973) 16:1207-1216. A linear correlation between the MR values and percent inhibition of p38α at 200 nM concentrations of the compounds was obtained. The smallest substituent (chloro) showed an MR value of about 6 and 52% inhibition at 200 nM; the compound with an intermediate MR value of about 8 (methoxy) provided 68% inhibition at this concentration, and the compound having the largest substituent (bromo) showed an MR value of approximately 9 and 72% inhibition at this concentration. Thus, it appears that the ability to inhibit p38α kinase is dependent on the size of the ortho substituent. While not intending to be bound by any theory, applicants believe that the bulkier substituents rotate the aroyl moiety out of the plane of the piperidine or piperazine ring.

20

Example 13


Additional Compounds

Additional compounds within the scope of the invention shown in Figures 1A-1I have been prepared.

Claims

1. The use of a compound of formula (3) to prepare a medicament for the treatment of conditions mediated by p38 α kinase, wherein the compound of formula (3) is of the formula:

5



wherein Ar¹ is a furanyl ring system optionally substituted by noninterfering substituents other than phenyl;

X¹ is CO or an isostere thereof;

Z is N or CH;

10

X² is CH₂ or an isostere thereof; and

Ph is optionally substituted phenyl.

2. The use of claim 1 wherein X¹ is CO and X² is CH₂.

15


3. The use of claim 1 or 2 wherein said furanyl ring system is optionally substituted by one substituent selected from the group consisting of alkyl, alkoxy, cycloalkyl, cycloheteroalkyl, heteroaryl, halogen, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, acyl, carboxy, hydroxy, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, CF₃, cyano, amino and alkylamino.

20

4. The use of any of claims 1-3 wherein Ph is optionally substituted by one substituent selected from the group consisting of alkyl, alkoxy, cycloalkyl, cycloheteroalkyl, heteroaryl, halogen, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, acyl, carboxy, hydroxy, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, CF₃, cyano, amino, alkylamino and aryl.

25

5. The use of a compound of formula (3) to prepare a medicament for treating conditions mediated by p38 α kinase wherein the compound of formula (3) is of the formula:



wherein Ar¹ is a thiophene ring system optionally substituted by a noninterfering substituent, and not fused to an additional ring;

X¹ is CO or an isostere thereof;

Z is N or CH;

5 X² is CH₂ or an isostere thereof; and


Ph is optionally substituted phenyl.

6. The use of claim 5 wherein X¹ is CO and X² is CH₂.

7. The use of claim 5 or 6 wherein said thiophene ring system is optionally substituted by at least one substituent selected from the group consisting of
10 alkyl, alkoxy, cycloalkyl, cycloheteroalkyl, heteroaryl, halogen, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, acyl, carboxy, hydroxy, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, CF₃, cyano, amino, alkylamino, and aryl.

8. The use of any of claims 5-7 wherein Ph is optionally substituted by one substituent selected from the group consisting of alkyl, alkoxy, cycloalkyl,
15 cycloheteroalkyl, heteroaryl, halogen, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, acyl, carboxy, hydroxy, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, CF₃, cyano, amino, alkylamino and aryl.

9. The use of a compound of formula (3) to prepare a medicament for the treatment of conditions mediated by p38α kinase wherein the compound of
20 formula (3) is of the formula:



wherein Ar¹ is an optionally substituted aryl group other than optionally substituted indolyl, benzimidazolyl, or benzotriazolyl;

X¹ is CO or an isostere thereof;

25 Z is N or CH;

X² is CH₂ or an isostere thereof; and

Ph represents optionally substituted phenyl

wherein at least one of Ar¹ and Ph is substituted by at least one substituent comprising a saturated ring system optionally containing 1-2 N atoms and/or 1-2 O atoms.

10. The use of claim 9 wherein X¹ is CO and X² is CH₂.

5 11. The use of claim 9 or 10 wherein Ar¹ is optionally substituted by at least one substituent selected from the group consisting of alkyl, alkoxy, cycloalkyl, cycloheteroalkyl, heteroaryl, halogen, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, acyl, carboxy, hydroxy, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, CF₃, cyano, amino, alkylamino, and aryl.

10 12. The use of any of claims 9-11 wherein Ph is optionally substituted by one substituent selected from the group consisting of alkyl, alkoxy, cycloalkyl, cycloheteroalkyl, heteroaryl, halogen, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, acyl, carboxy, hydroxy, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, CF₃, cyano, amino, alkylamino and aryl.

15 13. The use of a compound of formula (3) to prepare a medicament for the treatment of conditions mediated by p38 α kinase wherein the compound of formula (3) is a compound illustrated in Figures 1A-1I.

1/20




FIG. 1A

2/20







122	
123	
124	
119	
120	
121	

FIG. 1B

3/20







128		125	
129		126	
130		127	

FIG. 1C

4/20




FIG. 1D

5/20




FIG. 1E

6/20







<p>141</p> 	<p>142</p> 	<p>143</p> 	<p>144</p> 	<p>145</p> 	<p>146</p> 	<p>CHIRAL</p>
--	--	---	--	--	---	---------------

FIG. 1F

7/20







150		151		152	
147	CHIRAL 	148		149	

FIG. 1G

8/20





156			157
153		154	155
			

FIG. 1H

9/20






160		
161		
162		
	158	
	159	

FIG. 11

10/20




FIG. 1J

11/20




FIG. 1K

12/20







177		174	
178		175	
179		176	

FIG. 1L

13/20




FIG. 1M

14/20




FIG. 1N

15/20







195		192	
196		193	
197		194	

FIG. 10

16/20







201		198	
202		199	
203		200	

FIG. 1P

17/20




FIG. 1Q

18/20







213	
214	
215	
210	
211	
212	

FIG. 1R

19/20







219	
220	
221	
216	
217	
218	

FIG. 1S

20/20






225	
226	
222	
223	
224	

FIG. 17

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 September 2001 (07.09.2001)

PCT

(10) International Publication Number
WO 01/64676 A3

(51) International Patent Classification⁷: **C07D 471/10**,
235/06, 211/16, 401/06, 215/48, 401/14, 471/04, 405/12,
277/52, 209/08, 401/12, 403/12, 513/04, 473/06, 487/04,
409/06, 295/18, 211/28, 211/26, 401/10, 215/22, 307/68,
A61K 31/497, A61P 29/00, A61K 31/4545, 31/495,
31/496, 31/4535

(21) International Application Number: **PCT/US01/06715**

(22) International Filing Date: 28 February 2001 (28.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/185,571 28 February 2000 (28.02.2000) US

(71) Applicant: **SCIOS, INC.** [US/US]: 820 West Maude Avenue, Sunnyvale, CA 94086 (US).

(72) Inventors: **GOEHRING, Richard, R.**: 6195 Potters Lane, Pipersville, PA 18947 (US). **MAVUNKEL, Babu, J.**: 1348 Avoset Terrace, Sunnyvale, CA 94087 (US). **LIU, David, Y.**: 201 Fern Avenue, Palo Alto, CA 94306 (US). **SCHREINER, George, F.**: 12774 Leander Drive, Los Altos Hills, CA 94022 (US). **LUEDTKE, Gregory**: 963 La Mesa Terrace, #9, Sunnyvale, CA 94086 (US). **LEWICKI, John, A.**: 308 Escobar Avenue, Los Gatos, CA 95030 (US).

(74) Agents: **MURASHIGE, Kate, H.** et al.: Morrison & Foerster LLP, 3811 Valley Centre Drive, Suite 500, San Diego, CA 92130-2332 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:
with international search report

(88) Date of publication of the international search report:
28 March 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: INHIBITORS OF p38- α KINASE



(57) Abstract: The invention is directed to prepare medicaments for treatment of and to methods to treat conditions mediated by kinase using compounds of the formula (3) wherein Ar¹ and Ph are limited to specific embodiments or wherein the compound of formula (3) is a compound set forth in Figures 1A-11.

WO 01/64676 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCI/US 01/06715

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D471/10 C07D235/06 C07D211/16 C07D401/06 C07D215/48
 C07D401/14 C07D471/04 C07D405/12 C07D277/52 C07D209/08
 C07D401/12 C07D403/12 C07D513/04 C07D473/06 C07D487/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
P, A	WO 00 71535 A (SCIOS INC.) 30 November 2000 (2000-11-30) page 1 -page 3, line 22 ---	1-13
P, A	WO 00 12074 A (SCIOS INC.) 9 March 2000 (2000-03-09) cited in the application page 1 -page 4, line 10 ---	1-13
P, A	WO 00 59904 A (SCIOS INC.) 12 October 2000 (2000-10-12) cited in the application page 1 -page 4, line 5 ---	1-13
Y	WO 99 61426 A (SCIOS INC.) 2 December 1999 (1999-12-02) cited in the application page 1 -page 4, line 7 ---	1-38
-/--		



Further documents are listed in the continuation of box C



Patent family members are listed in annex

* Special categories of cited documents

A document defining the general state of the art which is not considered to be of particular relevance

L earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

I later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

Z document member of the same patent family

Date of the actual completion of the international search

18 December 2001

Date of mailing of the international search report

02/01/2002

Name and mailing address of the ISA

European Patent Office, P.O. Box 5818 Patentlaan 2
 NL 2280 HV Rijswijk
 Tel (+31-70) 340-2000, Tx 31 651 eponl
 Fax (+31-70) 340-3016

Authorized officer

Kyriakakou, G

INTERNATIONAL SEARCH REPORT

International Application No.

PCI/US 01/06715

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D409/06 C07D295/18 C07D211/28 C07D211/26 C07D401/10
C07D215/22 C07D307/68 A61K31/497 A61P29/00 A61K31/4545
A61K31/495 A61K31/496 A61K31/4535

According to International Patent Classification (IPC) or to both national classification and IPC.

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 28292 A (SMITHKLINE BEECHAM) 2 July 1998 (1998-07-02) cited in the application page 1 -page 5, line 5	1-38
Y	WO 98 06715 A (SMITHKLINE BEECHAM CORPORATION) 19 February 1998 (1998-02-19) cited in the application page 1 -page 5, line 15	1-13

☐ Further documents are listed in the continuation of part C.

☒ Patent family members are listed in annex

* Special categories of cited documents

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claims for which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *I* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *S* document member of the same patent family

Date of the actual completion of the international search

18 December 2001

Date of making of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5618 Patentlaan 2
NL 2280 HV Rijswijk
Tel (+31-70) 340 2046 Telex 51651 Epe nl
Fax (+31-70) 340 3016

Authorized officer

Kyriakakou, G

INTERNATIONAL SEARCH REPORT

Information on patent family members

In' tional Application No

PC1/US 01/06715

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 0071535	A	30-11-2000	AU	3920900 A		23-10-2000
			AU	5442400 A		12-12-2000
			WO	0059904 A2		12-10-2000
			WO	0071535 A1		30-11-2000
WO 0012074	A	09-03-2000	AU	5793699 A		21-03-2000
			BR	9913654 A		27-11-2001
			CN	1323209 T		21-11-2001
			EP	1107758 A2		20-06-2001
			WO	0012074 A2		09-03-2000
WO 0059904	A	12-10-2000	AU	3920900 A		23-10-2000
			AU	4092099 A		13-12-1999
			BR	9911069 A		06-02-2001
			CN	1305464 T		25-07-2001
			EP	1080078 A1		07-03-2001
			NO	20005881 A		09-01-2001
			WO	0059904 A2		12-10-2000
			WO	9961426 A1		02-12-1999
			AU	5442400 A		12-12-2000
			WO	0071535 A1		30-11-2000
WO 9961426	A	02-12-1999	US	6130235 A		10-10-2000
			AU	4092099 A		13-12-1999
			BR	9911069 A		06-02-2001
			CN	1305464 T		25-07-2001
			EP	1080078 A1		07-03-2001
			NO	20005881 A		09-01-2001
			WO	9961426 A1		02-12-1999
			AU	3920900 A		23-10-2000
			WO	0059904 A2		12-10-2000
WO 9828292	A	02-07-1998	WO	9828292 A1		02-07-1998
WO 9806715	A	19-02-1998	EP	0922042 A1		16-06-1999
			JP	2001506230 T		15-05-2001
			WO	9806715 A1		19-02-1998